

Randy Clark, MD
Revere Health Coral Desert Orthopedics
2825 south mall drive
St. George, UT 84790
435-628-9393 phone
435-628-9382 fax

The following applies to: Scar lysis and manipulation under anesthesia via
Knee arthroscopy or **Shoulder** arthroscopy,

Shoulder-Sling: For comfort as needed, may discontinue when pain is minimal.

Knee-crutches: May weight-bear as tolerated and discontinue crutches as able.

Physical therapy: Start formal physical therapy immediately and do the below listed exercises several times daily.

WOUND CARE:

- Shower: as long as bandages are covered with waterproof covering. You may remove the sling for showering.
- Bandages: Can remove 2-3 days after surgery, leave white strips in place.
- After bandages have been removed may shower without covering incision sites, keep out of direct spray from water and pat dry.
- NO soaking in bathtub/hot tub or pool for first 2-3 weeks.
- Signs of infection: excessive or worsening pain, increased redness around the incision sites, drainage from incisions that looks like pus and/or is foul smelling, if you run a fever above 101° F - **CALL IMMEDIATELY 435-628-9393**
- **PAIN/SWELLING CONTROL**
- Start your **pain medication** before your block wears off if you have one.
- For the first few days take your pain medication regularly as instructed, then wean as tolerated.
- **Ice** for 20-30 minutes at least four times per day for several days. You may use it more frequently as needed. It will help in reducing pain. Avoid placing ice directly on bare skin to prevent causing frostbite and give a break of 30 minutes in between sessions.
- It will help if you **sleep** with pillows behind your back and shoulder. Some patients find it helpful to sleep in a recliner.
- When dressing, put your operative arm in the sleeve first. When getting undressed, take your operative arm out last. Loose fitting, button-down shirts are recommended.

Post operative **knee** exercises (may start 1st day after surgery)

- You will begin flexion (bending) after surgery. Work on active bending (using your own muscle power) as far as possible and as tolerated. You can straighten using your good leg to help straighten the operated leg.
- Active control of quadriceps muscle
 - Exercises to be performed 2-3 times daily:
 - **Straight leg raises** while wearing brace locked straight, foot straight and turned out, lift leg up off bed/floor. Goal is 50-100 reps three times per day.
 - **Isometric quadriceps contractions** (letting the leg rest straight), squeeze the quad muscles without lifting the leg off the bed/floor.

(see page 2)

The following applies to: Scar lysis and manipulation under anesthesia via
Knee arthroscopy or **Shoulder** arthroscopy,

Post operative **shoulder** exercises (may start 1st day after surgery)

Elbow Range of Motion

1. Remove sling.
2. With elbow bent at 90°, bend and straighten elbow as though brushing something off the front of you. Repeat 30 times, 3 times per day.

Grip Strengthening

1. Make tight fist or grip rubber ball.
2. Hold for 10 seconds. Repeat 30 times, 3 times per day.

Scapula Retraction

1. Stand with arms at side.
2. Pinch shoulder blades together as if you are trying to get them to touch in the middle.
3. Hold for 10 seconds. Repeat 30 times, 3 times per day.

*The following is to be done without using your shoulder muscles but with the assistance of the non-operative hand to make the motion, it is described as if the right shoulder has been operated on.

Supported Arm Pendulums

1. Bend over at waist. Bend Right elbow 90° and support with Left hand.
2. Using your left arm to move the right, make small circles clockwise and counterclockwise. Repeat 30 times each way, 3 times per day.